GC/Chlamydia Sample/Order Requirements

Source	Tube	Test to Order
Vaginal With orders for both Vaginitis Panel and GC/Chlamydia	HOLOGIC 216 ml Aptima- Mainseet Study: Theresory Media (STM)	GC CHLAMYDIA PCR RRMC and VAGINITS PANEL RRMC NOTE: For patients younger than 14 order: GC/CHLAMYDIA AND VAGINITIS PANEL UVMC
Endocervical/Cervical Urethral	HOLOGIC 2.9 et. Aptima beet Spectruet Table	GC & CHLAMYDIA AMPLIFIED RNA MAYO
Urine	BECK/SHOOL Constant Fred Media Constant Fred Media State of the State of the Sta	GC CHLAMYDIA URINE AMPLIFIED RNA UVMC
Orophrayngeal (Throat) Anorectal (Anus/Rectum)	HOLOGIC 216 mi Aptima Millioner Swith Transport Minita (STM)	GC CHLAMYDIA AMPLIFIED RNA (SWAB) UVMC
Occular	HOLOGIC 29 ml Aptima • Minimer Swith Threesery Minima (19 ml OC HOLOGIC 29 mL Aptima Bent Spectrum Taserie Tabe	GC CHLAMYDIA MISC SITES (OCULAR) MAYO

A source must ALWAYS be entered for GC/Chlamydia tests. N/A IS NOT acceptable. The reference lab WILL NOT run the test without source information.

If a source is not indicated on the sample or the req, call the provider to find out.